

4x Vaccinated Youth 318% More Likely to Die Than Unvaxxed Peers

Analysis by [Dr. Joseph Mercola](#)

April 13, 2024

STORY AT-A-GLANCE

- › Young people who received multiple COVID jabs were significantly more likely to die than those who skipped the shots, according to data from the U.K.'s Office for National Statistics
- › An analysis by The Exposé revealed that, in February 2023, those who received four COVID-19 shots had a 318% higher mortality rate than the unvaccinated group
- › The other months analyzed showed four-shot teens and youth were between 221% and 290% more likely to die than those who didn't get the shot
- › Another study found that, compared to unvaccinated children, vaccinated children had significantly higher rates of asthma, allergies, eczema, respiratory infections, behavioral issues and other health conditions
- › Separate research showed "for every life saved, there were nearly 14 times more deaths caused by the modified mRNA [COVID-19] injections"

Young people who received multiple COVID jabs were significantly more likely to die than those who skipped the shots, according to data from the UK's Office for National Statistics (ONS).^{1,2} The data include deaths by vaccination status from April 1, 2021, to May 31, 2023, when the COVID-19 shot campaign was in full effect.

When The Exposé analyzed the data, they revealed the disturbing finding that those with the most COVID-19 shots fared the worst:³

"Our analysis focused on mortality rates per 100,000 person-years from January to May 2023 among residents in England aged 18 to 39, and what we found is truly shocking. Initial observations of the data prove that individuals in this age bracket who had received four doses of a COVID-19 vaccine exhibited higher mortality rates compared to their unvaccinated counterparts."

Mortality Rate 318% Higher Among Quadruple-Jabbed Youth

As noted by Canadian oncologist and cancer researcher Dr. William Makis, "The more mRNA shots you take, the greater the immune system damage," which can lead to a host of health problems, not the least of which is a greater risk of impaired cancer surveillance and **turbo cancer**.⁴

The Exposé analysis suggests multiple shots also raise mortality rates among those aged 18 to 39 years. In every month, those who received four COVID-19 shots were significantly more likely to die than those who hadn't received any. In January 2023, for instance, those who never received a COVID-19 shot had a mortality rate of 31.1 per 100,000 person-years.

But among the quadruple jabbed, the rate was 106 per 100,000 person-years. Even among those who had received one COVID-19 shot, the mortality rate was much higher than the unvaccinated at 53.3 per 100,000 person-years that month. According to The Exposé:⁵

"For the remaining months, unvaccinated teens and young adults mortality rate remained within the 20-something per 100,000 person-years. Whereas four-dose vaccinated teens and young adults' mortality rates only went as low as 80.9 per 100,00 in April and remained within 85 to 106 per 100,000 for the remaining months.

The January to May average mortality rate per 100,000 person-years was 26.56 for unvaccinated teens and young adults and a shocking 94.58 per 100,000 for four-dose vaccinated teens and young adults. Meaning on average, the four-

dose vaccinated were 256% more likely to die than the unvaccinated based on mortality rates per 100,000."

In February 2023, however, those who received four COVID-19 shots had a 318% higher mortality rate than the unvaccinated group. The other months showed four-shot teens and youth were between 221% and 290% more likely to die than those who didn't get the shot.⁶

"These figures are extremely worrying and strongly suggest that Covid-19 vaccination increases a person's mortality rate, which in turn suggests Covid-19 vaccination may actually be killing teens and young adults in the tens of thousands," The Exposé reported.⁷

COVID-19 Shots Killed More People Than They Saved

Other research has reached similar conclusions that mRNA COVID shots may be deadly. A now-retracted narrative review published in the journal *Cureus* called for a global moratorium on mRNA COVID-19 shots,⁸ citing significant increases in serious adverse events among those who received the injections, along with an "unacceptably high harm-to-reward ratio."⁹

When factoring in absolute risk and the "number needed to vaccinate" (NNV), a metric used to quantify how many people need to be vaccinated to prevent one additional case of a specific disease, the review found "for every life saved, there were nearly 14 times more deaths caused by the modified mRNA injections."¹⁰

As for why the paper was retracted, study author Steve Kirsch said, "It's about supporting the narrative."¹¹ Board-certified internist and cardiologist Dr. Peter McCullough, another of the paper's authors, called the retraction a "stunning act of scientific censorship."¹²

In addition to calling for a global moratorium on mRNA COVID-19 shots, the authors of the paper — M. Nathaniel Mead, Stephanie Seneff, Ph.D., Russ Wolfinger, Ph.D., Jessica Rose, Ph.D. Kris Denhaerynck, Ph.D., Kirsch and McCullough — said the shots should be

immediately removed from the childhood vaccine schedule, while boosters should also be suspended.

"It is unethical and unconscionable to administer an experimental vaccine to a child who has a near-zero risk of dying from Covid-19 but a well-established 2.2 percent risk of permanent heart damage based on the best prospective data available," the paper notes.¹³

The moratorium is warranted based on the shots' risks of serious adverse events, the mechanisms behind those adverse events, mortality data and issues with inefficacy, vaccine control and processing.¹⁴ According to the review:¹⁵

"Federal agency approval of the COVID-19 mRNA vaccines on a blanket-coverage population-wide basis had no support from an honest assessment of all relevant registrational data and commensurate consideration of risks versus benefits.

Given the extensive, well-documented SAEs [serious adverse events] and unacceptably high harm-to-reward ratio, we urge governments to endorse a global moratorium on the modified mRNA products until all relevant questions pertaining to causality, residual DNA, and aberrant protein production are answered."

Are Unvaxxed Children Healthier Than Vaxxed?

Another study that runs counter to the narrative was also retracted. The research, by Dr. Paul Thomas, who had his medical license suspended after he became widely known for supporting informed consent for vaccinations, and James Lyons-Weiler, with The Institute for Pure and Applied Knowledge (IPAK), analyzed data spanning a 10-year period within a pediatric practice.

It focused on comparing health outcomes between vaccinated and unvaccinated children.¹⁶ The data showed vaccinated children had significantly higher rates of:¹⁷

Asthma	Allergies
Eczema	Sinusitis
Gastroenteritis	Respiratory infections
Middle ear infection	Conjunctivitis
Breathing issues	Behavioral issues

Further, none of the 561 children who were not vaccinated had attention deficit hyperactivity disorder (ADHD), while 0.063% of partially and fully vaccinated children did. "The implications of these results for the net public health effects of whole-population vaccination and with respect for informed consent on human health are compelling," they wrote.¹⁸

The study also points out that the rate of autism spectrum disorder in their practice was half that of the U.S. national average (0.84% versus 1.69%). The rate of ADHD in the practice was also about half the national rate. According to the authors, "The data indicate that unvaccinated children in the practice are not unhealthier than the vaccinated and indeed the overall results may indicate that the unvaccinated pediatric patients in this practice are healthier overall than the vaccinated."¹⁹

The researchers call for more studies on this topic to be done independently from the vaccine industry, emphasizing the need for unbiased research in understanding the impact of vaccinations on children's health. Canadian oncologist and cancer researcher Dr. William Makis explained:²⁰

"The results were not even remotely close in comparison. It is a total knockout. If you listened to the TV Media you would think that results would be just the opposite of these findings. There is a reason in 100 years of Vaccinating Children there have NEVER been retrospective studies.

Don't you want to see 5-, 10-, 15- or even 20-year studies of the vaxxed vs unvaxxed? Wouldn't that be VERY Helpful? Why has there never been any studies like this?

They would 100% PROVE without Shadow of a Doubt that not only do Vaccines NOT prevent infections and transmission but they are severely harming children – especially for Allergies, Middle Ear Infections, Autism, Breathing issues, Sinusitis, Respiratory Infections, Eye Infections, Gastroenteritis, Eczema and Behavioral issues.

IF Vaccines PREVENT Infections then why are infections Significantly Higher in all of Dr. Thomas' vaccinated children? These are serious life changing chronic conditions. Vaccines shift the immune system to Allergy and Autoimmunity, and you have more and more infections of other kinds.

In short, what does this mean? It means that Vaccines DESTROY a Child's Immune System, every vax is different with different live viruses and toxins and they will really hit a child with a weaker immune system first."

Other Studies Show Health Issues More Common in Vaxxed Children

While the mainstream narrative only shares the notion that "vaccines are safe and effective," several studies have shown this isn't always the case. One study looked at health outcomes of **vaccinated and unvaccinated children** from three medical practices in the U.S.²¹

Vaccinated children were much more likely to have several health issues compared to unvaccinated children. Specifically, the chances of being diagnosed with severe allergies, autism, gastrointestinal disorders, asthma, ADHD and chronic ear infections were significantly higher for vaccinated children.

For example, vaccinated children had over four times the risk of severe allergies and over 20 times the risk of ADHD compared to unvaccinated children. The vaccinated children who were not breastfed or delivered by cesarean section had the highest risk of negative health outcomes.

Another study found that children vaccinated before their first birthday had a higher chance of experiencing developmental delays, asthma and ear infections.²² The more vaccine doses the children received, the higher their risk for these health issues became. Further, when the scientists looked at developmental delays over time, the risk increased as children got older, from 6 months up to 24 months of age.

The risk for four health conditions – developmental delays, asthma, ear infections and gastrointestinal disorders – also increased when they extended the age for possible diagnosis from at least 3 years to at least 5 years. In another example, researchers looked into the health of U.S. children who are homeschooled, comparing those who had been vaccinated with those who hadn't.²³

While the vaccinated children were less likely to have had chickenpox or whooping cough (pertussis), they were more likely to have had pneumonia, ear infections, allergies, and neurodevelopmental disorders such as a learning disability, ADHD or autism.

A particularly high risk of neurodevelopmental disorders (NDDs) was found in children who were both born prematurely and vaccinated, with these children being 6.6 times more likely to have NDDs compared to other children.

The fact that no link was found between premature birth and NDDs among the unvaccinated raises the disturbing possibility that the vaccination schedule for premature babies could be responsible for the neurological disorders some premature babies exhibit, which have been previously assumed to be simply a result of premature birth.

Past research published in the journal *Human & Experimental Toxicology* also showed infant mortality rates correlated with childhood vaccination rates, with high-uptake

countries having higher child mortality.²⁴ A reanalysis of the study, published in the peer-reviewed journal *Cureus* in February 2023, reaffirmed the positive correlation between number of **vaccine doses and infant mortality rates**.²⁵

As it stands, the childhood vaccine schedule shouldn't be considered safe and effective for all – and mRNA COVID-19 shots' association with increased mortality and other significant health issues should give everyone pause. If you've developed unusual symptoms after a COVID-19 shot or other vaccination, seek out help from an expert.

The Front Line COVID-19 Critical Care Alliance (FLCCC) also has a treatment protocol for COVID-19 shot injuries. It's called **I-RECOVER** and can be downloaded from covid19criticalcare.com.²⁶

Sources and References

- ^{1, 3, 5, 6, 7} [The Expose March 3, 2024](#)
- ² [U.K. Office for National Statistics, Deaths by vaccination status, England, August 25, 2023](#)
- ⁴ [Rumble, The HighWire with Del Bigtree, Is the COVID Vaccine Causing Turbo Cancers? September 22, 2023](#)
- ⁸ [Cureus January 24, 2024](#)
- ^{9, 15} [Cureus January 24, 2024, Abstract](#)
- ¹⁰ [Cureus January 24, 2024, Review](#)
- ¹¹ [Substack, Steve Kirsch's newsletter February 18, 2024](#)
- ¹² [Substack, Courageous Discourse February 24, 2024](#)
- ¹³ [World Tribune February 5, 2024](#)
- ¹⁴ [Children's Health Defense January 29, 2024](#)
- ^{16, 18, 19} [Int J Environ Res Public Health. 2020 Nov; 17\(22\): 8674](#)
- ^{17, 20} [Substack, COVID Intel, Dr. William Makis April 2, 2024](#)
- ²¹ [JTranslSci 7: June 12, 2021 DOI: 10.15761/JTS.1000459](#)
- ²² [SAGE Open Med. 2020; 8: 2050312120925344](#)
- ²³ [J Transl Sci 3: April 24, 2017 DOI: 10.15761/JTS.1000186](#)
- ²⁴ [Human & Experimental Toxicology May 4, 2011](#)
- ²⁵ [Cureus February 2, 2023; 15\(2\): e34566](#)
- ²⁶ [Covid19criticalcare.com](#)