MERCOLA
Take Control of Your Health

How Strength Training Supports Cardiovascular Health

Analysis by Dr. Joseph Mercola | November 06, 2026

STORY AT-A-GLANCE

- Strength training supports cardiovascular health alongside muscle and bone health, with research showing improvements in blood pressure, resting heart rate, blood vessel function and autonomic regulation of the heart
- Progressive resistance programs typically involved two to three sessions per week for eight to 24 weeks, giving you a practical framework for building cardiovascular benefits alongside strength
- Gradually increasing resistance as you become stronger helps provide the continued training stimulus your heart, arteries and muscles need to adapt
- Blood-flow restriction (BFR) training, such as the branded KAATSU system, provides a lower-load strength-training option if you're frail, deconditioned, recovering from an injury or unable to tolerate conventional heavy lifting. This approach is separate from the cited research and wasn't evaluated in the reviewed studies
- Strength training works best as part of a broader heart-health strategy that includes daily walking, frequent movement instead of prolonged sitting and limiting linoleic acid from seed oils

Most people think of strength training as a way to build muscle and protect bone. But a growing body of research shows it also reshapes your cardiovascular system — improving blood pressure, arterial flexibility and the nervous system's regulation of your heart.

A 2026 systematic review and meta-analysis published in the Journal of Applied Physiology examined 60 studies and found evidence of these benefits in postmenopausal women, a population at elevated cardiovascular risk.[1] The strongest effects appeared in systolic blood pressure and mean blood pressure.

Those findings matter because cardiovascular risk rises after menopause, while changes in blood vessel function and autonomic control — the part of your nervous system that regulates heart rate and blood pressure without conscious effort — place additional strain on your heart and circulation.

Systolic blood pressure is the top number in a blood pressure reading and reflects the pressure in your arteries when your heart contracts. Mean blood pressure represents the average pressure your arteries experience throughout each heartbeat. Both provide important clues about the workload placed on your cardiovascular system.

The review also documented smaller improvements in diastolic blood pressure — the bottom number, reflecting pressure between heartbeats — and resting heart rate, though these effects were modest compared with the changes in systolic and mean blood pressure.

Beyond blood pressure, the researchers found modest improvements in vascular function and heart rate variability. Heart rate variability measures the differences in timing between heartbeats and provides insight into how well your autonomic nervous system regulates your heart.

The review also found no adverse structural changes to the heart in the small number of studies (three) that assessed this outcome — addressing a longstanding concern that heavy resistance exercise might thicken the heart wall in unhealthy ways — which suggests strength training may give you a practical way to target muscle, bone and cardiovascular health with the same form of exercise.

The question, then, isn't simply whether lifting weights makes you stronger. It's what repeated resistance does to the heart, arteries and nervous system over time. The findings reveal several ways strength training changes cardiovascular function from the

inside out.

Progressive Strength Training Improves How Your Heart and Blood Vessels Work

The studies included 1,931 postmenopausal women ages 52 to 78. Some were healthy women while others had uncomplicated **high blood pressure**, giving researchers a broad look at how resistance training affects cardiovascular function after **menopause**.[2]

The researchers isolated **resistance training** rather than combining it with aerobic exercise. That distinction helps show what strength work itself contributes. The interventions lasted at least four weeks, although programs that assessed cardiovascular outcomes commonly ran eight to 24 weeks.

The review's authors also noted that most included trials carried a higher risk of bias and did not evaluate hard clinical outcomes such as heart attacks, strokes or cardiovascular mortality — the findings described here reflect improvements in blood pressure and vascular markers, not confirmed reductions in cardiovascular events.

- **Most successful programs followed a practical, progressive structure —** Participants generally trained two or three times per week, completing both upper- and lower-body exercises such as leg presses, chest presses, rows and leg curls. Most programs prescribed two or three sets of eight to 15 repetitions, with resistance equal to about 50% to 80% of the maximum weight a participant could lift once. Sessions generally lasted 40 to 60 minutes.

  Progress mattered as much as participation. Loads typically rose about 2% to 10% as participants completed more repetitions, reported less effort or hit their target sets with less strain. Repeating the same easy workout indefinitely stops producing cardiovascular adaptation — your body needs a gradually greater challenge to keep improving.

- **Greater training doses were linked to stronger cardiovascular responses —** The researchers noted that frequency, total exercise volume and program duration helped explain why cardiovascular improvements differed among studies. Higher weekly exercise doses were associated with greater improvements in hemodynamics — essentially, how efficiently blood moves through your body and how much strain that movement places on your heart and arteries.

  Shorter programs or those without adequate progression appeared less likely to provide enough stimulus for meaningful cardiovascular or blood vessel adaptation. This makes consistency a useful target you can track: complete your planned sessions, record your loads and gradually build from there.

- **Blood vessels became more flexible and responsive —** Thirteen studies examined vascular structure or function. Some reported better endothelial function, improved blood flow, lower resistance in peripheral blood vessels and greater nitric oxide availability. The endothelium is the thin inner lining of your blood vessels, while nitric oxide helps those vessels relax and widen so blood flows more freely.

  Several studies also reported decreases of about 0.6 to 0.8 meters per second in pulse wave velocity, a measure of how quickly pressure waves move through your arteries. Slower pulse wave velocity generally reflects less arterial stiffness. A useful comparison: pressure waves travel faster through a rigid pipe than through a flexible hose. Slower velocity tells you your arteries are staying supple.

- **Women with greater vascular problems appeared to have more room to improve —** The review noted that reductions in arterial stiffness tended to be greater in people who started with higher stiffness or high blood pressure. That's an important practical point: resistance training isn't reserved for people who already have ideal cardiovascular health.

  At the same time, vascular results differed across studies, and fewer trials examined these outcomes than blood pressure. The researchers therefore cautioned against treating one specific resistance-training program as the definitive

prescription for every woman.

- **Strength training changes both blood vessels and nervous system control —** Repeated muscle contractions alter blood flow and create shear stress, the gentle friction of blood against vessel walls. This stimulus supports greater nitric oxide availability, less oxidative stress and better vessel dilation. Resistance training also appears to improve the flexibility of vascular smooth muscle and remodel smaller arteries, reducing resistance against which your heart must pump.

  Nervous system changes add another layer. Studies of **heart rate variability** showed shifts toward stronger parasympathetic activity — the calming "rest-and-recover" side of your autonomic nervous system. Together, these vascular and nervous system adaptations help explain why progressive resistance exercise produces benefits that reach well beyond stronger muscles.

Build Cardiovascular Resilience with Progressive Strength Training

These findings give you a clear rationale for adding structured resistance exercise to your weekly routine — and for progressing it over time rather than staying at the same easy level. But strength training works best as one component of a broader strategy.

Your heart, blood vessels, muscles and metabolism all respond to what you do every day, so I'd address the major drivers together rather than rely on one form of exercise alone. Progressive resistance training gives your cardiovascular system an important physical stimulus, while regular walking, lower **linoleic acid** (LA) intake and steady daily movement support heart health from different angles.

1. **Make strength training a consistent part of your week —** Aim for two to three resistance-training sessions each week. That frequency closely matches the programs examined in the research, which commonly lasted eight to 24 weeks.[3] If you're starting from a low activity level, consistency matters more than chasing heavy weights. Build the habit first, then progress.

If you're unsure how to estimate your starting weight, choose a load that lets you complete your target repetitions with good form while the last two or three feel genuinely challenging. That typically lands you in the right intensity range without formal testing.

2. **Increase resistance gradually as your body adapts —** Progressive overload means slowly asking your muscles to do more. In the studies, resistance commonly increased by about 2% to 10% based on improved performance, completed repetitions or perceived effort. Track your weights and repetitions in a simple log so progress is visible rather than assumed.

3. **Consider BFR training if conventional strength training is too demanding —** If you're frail, deconditioned or recovering from an injury, BFR training — sold under branded systems such as KAATSU — gives you another way to stimulate your muscles with much lighter loads. BFR uses specialized cuffs to partially restrict blood flow during exercise, allowing low-intensity movements to create a stronger muscular training signal.

   Note that BFR training was not part of the cited resistance-training review; this is a separate option worth discussing with a qualified trainer or healthcare provider when heavy lifting isn't appropriate or tolerable.

4. **Walk every day and break up long periods of sitting —** Strength training is only one aspect of cardiovascular health. Your heart also benefits from frequent, lower-intensity movement throughout the day, which is why I recommend working toward about 60 minutes of walking daily. If that's more than you're doing now, start where you are and add five to 10 minutes per week.

   If you spend much of your day seated, stand up and move at regular intervals as well. Walking provides a steady cardiovascular stimulus, while resistance exercise challenges your body in a different way. Together, they create a more complete movement strategy.

5. **Lower your LA intake to support cardiovascular and mitochondrial health —** Strength training improves how your blood vessels function, but those vessels depend on healthy mitochondria to sustain that function over time. That's why it's important to pair strength training and daily walking with another foundational step: reducing excess LA, the primary polyunsaturated fat found in **seed oils**. I recommend keeping LA below 5 grams per day, and ideally closer to 2 grams.

   The biggest sources include soybean, corn, sunflower, safflower, cottonseed and canola oils. Excess LA becomes incorporated into mitochondrial membranes, where it oxidizes into damaging compounds that interfere with cellular energy production. Reducing these oils while emphasizing fats such as grass fed butter, ghee or tallow addresses cardiovascular health from another direction, by supporting the cellular energy systems your heart and blood vessels depend on.

FAQs About Strength Training for Cardiovascular Health

Q: How does strength training support cardiovascular health?

A: Resistance training does more than increase muscle strength. It supports healthier blood pressure, more responsive blood vessels and better regulation of your heart by the autonomic nervous system. Research in postmenopausal women also found no harmful structural changes to the heart from resistance training.

Q: How often do I need to strength train to get cardiovascular benefits?

A: Most of the programs studied involved two to three strength-training sessions per week and commonly lasted eight to 24 weeks. Gradual progression matters, too. As you get stronger, increasing the resistance gives your body a continued stimulus to adapt rather than allowing your workouts to become too easy.

Q: Does strength training improve the health of your arteries?

A: Yes. Studies found that blood flow improved, nitric oxide availability increased and several other markers of vascular function shifted in a favorable direction. Some also found reductions in arterial stiffness, including decreases of roughly 0.6 to 0.8 meters per second in pulse wave velocity, which measures how quickly pressure waves travel through your arteries.

Q: What if I'm too frail or injured to lift heavy weights?

A: BFR training, including branded systems such as KAATSU, offers an alternative when conventional heavy resistance training isn't appropriate or tolerable. Specialized cuffs partially restrict blood flow while you perform low-intensity exercise, allowing lighter movements to provide a stronger muscular training stimulus. This approach was not evaluated in the cardiovascular research discussed above.

Q: Is strength training enough to protect my cardiovascular health?

A: No. I recommend treating resistance exercise as one part of a broader strategy. Work toward about 60 minutes of walking each day, interrupt prolonged sitting with regular movement and reduce your intake of LA from seed oils such as soybean, corn, sunflower, safflower, cottonseed and canola oils. These steps support cardiovascular health through complementary pathways rather than relying on exercise alone.

This article is for informational purposes only and does not constitute medical advice. Consult a qualified healthcare provider before making changes to your health regimen.

Sources and References

- [1, 2, 3] Journal of Applied Physiology 2026, Volume 141, Issue 2