

Antioxidant Vitamins and Zinc Slowed Vision Loss in Advanced Age-Related Macular Degeneration

Analysis by [Dr. Joseph Mercola](#)

October 04, 2026

STORY AT-A-GLANCE

- › In the Age-Related Eye Disease Study (AREDS), a randomized clinical trial, a specific combination of antioxidant vitamins and zinc lowered the risk of progressing to advanced age-related macular degeneration (AMD) and helped preserve vision in people who already had intermediate disease or advanced disease in one eye, but it did not prevent the condition in healthy eyes
- › Your retina contains some of the highest concentrations of zinc in your body – and research indicates this mineral supports cell repair, communication between retinal cells, and the enzymes that help protect eye tissue against everyday damage
- › Maintaining the right zinc balance is just as important as getting enough because zinc deficiency impairs retinal function. Laboratory studies indicate that excessive zinc concentrations can also damage retinal cells
- › Animal foods such as oysters, grass fed beef, and lamb provide zinc in a form your body absorbs more efficiently than many plant foods, making animal-based whole-food sources an important part of supporting long-term eye health
- › If you eat large amounts of whole grains or legumes, remember that they contain natural compounds called phytates that reduce zinc absorption. Eating zinc-rich animal foods as part of a balanced meal helps improve the amount your body actually uses

Your eyes rely on a constant supply of nutrients to preserve the sharp central vision you need to read, drive, recognize faces, and perform countless everyday tasks. Many people assume a healthy diet covers those needs, yet a landmark clinical trial sponsored by the National Eye Institute found that for people already at risk of severe vision loss, a targeted nutrient formula measurably slowed disease progression.¹

The nutrient that emerged as especially important is zinc, a mineral your body uses in hundreds of biological processes every day and one your retina demands in unusually large quantities. What makes zinc unusual isn't that your eyes need it. It's that your eyes are remarkably sensitive to how much of it they get.

That tension between enough and too much runs through every layer of the science covered in this article, from the clinical trial that changed how specialists recommend nutritional support for age-related macular degeneration (AMD) to laboratory discoveries about what happens inside retinal cells when this mineral falls out of balance. Your retina is especially vulnerable to oxidative stress — the accumulation of unstable molecules produced during normal metabolism that, left unchecked, damage cells from the inside out.

Antioxidant Vitamins and Zinc Slowed Vision Loss in People at Highest Risk

The Age-Related Eye Disease Study (AREDS) was designed to determine whether a specific combination of antioxidant vitamins and zinc could slow the progression of **AMD** in people at risk of severe vision loss.^{2,3} The randomized clinical trial enrolled 4,757 adults between the ages of 55 and 80 at 11 clinical centers across the U.S. Of those, 3,640 were randomized to the AMD treatment arms and followed for an average of 6.3 years.

Participants received one of four treatments: **zinc** alone, antioxidant vitamins alone, antioxidants plus zinc or a placebo. Rather than testing whether supplements improved healthy eyes, the researchers focused on slowing disease progression in people with

varying stages of AMD.

- **The strongest results occurred in participants who already had more advanced disease** — The study found that the greatest benefit occurred in people with intermediate AMD or advanced AMD in one eye. For this group, the combination of antioxidants and zinc reduced the risk of progressing to advanced AMD by about 25% and lowered the risk of vision loss by about 19%.

Those percentages are relative, meaning they describe how much lower the risk was in the treated group compared with the placebo group — not a 25% chance of holding on to your vision. They also apply only to people who were already at high risk when the study began.*

In contrast, participants with no AMD or only early disease didn't experience the same measurable benefit. These findings positioned the AREDS formula as a targeted intervention for people at highest risk, not a preventive supplement for the general population.

- **Combining antioxidants with zinc worked better than using either treatment alone** — Researchers compared four treatment strategies and found a clear pattern. Zinc on its own did show a benefit, but only in the group at highest risk — and even there, the finding was borderline. Antioxidant vitamins, on their own, never produced a result the researchers could separate from chance.

The combination of antioxidants and zinc consistently produced the greatest benefit, outperforming either approach by itself.

Participants receiving the combined formula had the lowest likelihood of progressing to advanced AMD, while those receiving the placebo had the highest risk. The results led the researchers to recommend the combination rather than either component alone. The trial was not designed to formally test how the nutrients interact with one another.

- **The researchers tested one precise formula instead of general vitamin use** — The benefits came from a carefully standardized combination consisting of 500 milligrams (mg) of vitamin C, 400 international units of vitamin E, 15 mg of beta-carotene, 80 mg of zinc as zinc oxide, and 2 mg of copper as cupric oxide.

That zinc dose is far above ordinary intake — the Recommended Dietary Allowance (RDA) is 11 mg daily for men and 8 mg for women, and the Tolerable Upper Intake Level for adults is 40 mg daily.⁴ It was a deliberately high research dose given under trial supervision, not a general intake recommendation.

Copper was included as a precaution, because high-dose zinc can interfere with copper absorption. In AREDS, annual blood sampling in a subset of participants showed no increased risk of copper-deficient anemia in those assigned to zinc.

The study also found that nearly two-thirds of participants already took a daily multivitamin, yet those at high risk still received additional protection from the AREDS formula beyond what a standard multivitamin provided. Because the multivitamin offered in the trial also contained 15 mg of zinc, some participants were taking as much as 95 mg of zinc daily.

- **The study also clarified what the formula didn't accomplish** — Although the nutrient combination slowed progression to advanced AMD, it didn't restore vision that had already been lost. The researchers also evaluated **cataracts** and found that participants taking zinc alone, antioxidants alone, or the combined formula developed cataracts at approximately the same rate as those taking the placebo.

These findings narrowed the role of the formula to one specific purpose — slowing progression to advanced AMD in people already at elevated risk.

- **Long-term treatment proved generally safe but still required important precautions** — Researchers monitored participants throughout the seven-year study for adverse effects and reported relatively few serious problems overall.

However, about 7.5% of participants receiving zinc-containing treatments experienced urinary tract conditions requiring hospitalization compared with 5% of those who didn't receive zinc. That difference was statistically significant, and it was larger among men considered alone (8.6% versus 4.4%).

Mild anemia also occurred somewhat more often in participants taking zinc, although laboratory testing didn't show meaningful differences between treatment groups. The researchers also highlighted an important safety concern regarding beta-carotene, noting its association with increased lung cancer risk in smokers. That finding eventually led to updated AREDS2 formulations that replaced beta-carotene with lutein and zeaxanthin.

In AREDS2, current smokers were excluded from beta-carotene randomization, and the excess lung cancers that were observed occurred mostly in former smokers (23 cases versus 11 cases).

- **A follow-up trial updated the original formula based on what the first study revealed** — The AREDS2 study, completed in 2013, made three key changes.⁵ Researchers replaced beta-carotene with 10 mg of **lutein** and 2 mg of zeaxanthin, two carotenoids that naturally concentrate in the macula. They also tested whether reducing the zinc dose from 80 mg to 25 mg affected the formula's effectiveness and found that the lower dose provided similar protection against progression to advanced AMD.

There was no statistically significant difference between the two doses, and the review authors note that this raises a reasonable question about whether the higher dose is still warranted.⁶

The study also evaluated omega-3 fatty acids but found they didn't provide additional benefit when added to the formula. Today, most eye health supplements sold as "AREDS" formulas are based on the AREDS2 formulation, making it the version most people are likely to encounter.

The AREDS trial indicated that a zinc-containing formula helped slow disease progression, but it didn't explain why the retina needs so much of it in the first place or what happens inside retinal cells when zinc levels drift too high or too low. A comprehensive review of decades of laboratory and clinical research offers those answers.

**These findings are from research conducted in clinical settings. Results may not apply to all individuals.*

Researchers Found That Healthy Eyes Depend on Balanced Zinc Levels

For a review published in Progress in Retinal and Eye Research, researchers reviewed decades of laboratory research, clinical trials and population studies to understand exactly how zinc supports healthy vision and why it's become one of the most important nutrients studied for eye health.⁷

The review summarized current evidence on how zinc functions inside the retina, how the body absorbs and regulates it, what happens when zinc levels become too low or too high and why researchers continue investigating its role in eye disease.

Unlike studies that focus only on people with AMD, this review also surveyed the literature on healthy eyes, inherited disorders, digestive diseases and other conditions that affect zinc status. The researchers concluded that adequate zinc is required for normal retinal function, while both deficiency and excess interfere with the cells responsible for healthy vision.

- **The retina naturally contains some of the highest zinc concentrations in the body**
— Researchers explained that the retinal pigment epithelium and choroid — the layers that nourish and support the retina — contain the greatest concentrations of zinc anywhere in the eye. These tissues do far more than store the mineral. Concentration is not the same as quantity, though.

The review's own accounting places the eye at under 0.01% of total body zinc, with most of the body's supply held in skeletal muscle and bone.

Zinc supports hundreds of enzymes along with protein production, cell repair, communication between retinal cells, and the activity of protective enzymes that defend eye tissue against everyday damage. The review is careful to note that zinc is not an antioxidant per se — it does not participate in redox reactions directly, but stabilizes and supports the enzymes that do.

The review also noted that scientists still don't fully understand every molecular pathway through which zinc protects retinal cells, making it one of the most actively studied nutrients in vision research.

- **Zinc deficiency affected eyesight long before permanent damage became obvious** — The review described several situations in which inadequate zinc disrupted normal vision. People with acrodermatitis enteropathica — a rare inherited disorder that impairs intestinal zinc absorption — develop eye abnormalities that are largely front-of-the-eye problems, including blepharitis, conjunctivitis, and corneal opacities.

In Crohn's disease, reduced retinal zinc has been implicated in a bilateral maculopathy with vision impairment that was responsive to zinc supplementation.

Long-term intravenous nutrition has likewise been associated with impaired visual function and low plasma zinc, though the review does not report reversal with zinc replacement in that group. These are rare, severe systemic deficiency states, not conditions the review extends to well-nourished people.

Researchers also reported that zinc deficiency often reduced the eye's ability to adjust from bright light to darkness, a process called dark adaptation. The clinical evidence here comes largely from people with alcoholism and hepatic cirrhosis, and the mechanism is indirect — zinc is a cofactor for the enzyme that converts vitamin A to retinal — so, these cases often require both zinc and vitamin A to improve.

When dark adaptation slows, driving at night or walking through dimly lit areas becomes much more difficult. Animal studies further showed that severe post-weaning zinc deficiency damaged photoreceptors — the specialized cells that detect light — along with the retinal pigment epithelium, although the same review notes that postnatal zinc deficiency on its own produced functional changes without visible structural damage.

The picture is not one-directional — in one rodent experiment, rats fed a low-zinc diet showed less light-induced retinal damage than those on regular or high-zinc diets.

- **More zinc didn't produce better results** — One of the review's clearest messages is that healthy vision depends on maintaining zinc within a normal range rather than consuming as much as possible. Laboratory studies showed that excessively high zinc concentrations triggered harmful chemical reactions inside retinal tissue by increasing reactive oxygen species, unstable molecules that damage proteins, fats, and DNA.

These were in vitro experiments on isolated retinal tissue exposed to zinc concentrations of 500 micromolar or more, which is far above anything produced by ordinary dietary intake, and not directly comparable to an oral supplement dose.

High zinc levels also reduced the activity of important antioxidant enzymes, weakening the eye's natural defense systems. To prevent both deficiency and overload, the body relies on specialized transport proteins that continually move zinc into and out of cells to maintain a healthy balance.

- **The foods you choose influence how much zinc your body actually uses** — Researchers emphasized that eating zinc and absorbing zinc are not the same thing. Animal foods such as red meat and shellfish provide zinc in a form your body absorbs more efficiently than many plant foods. In contrast, phytates — compounds naturally present in whole grains and legumes — bind to zinc in the digestive tract and reduce the amount your body absorbs.

The review also found that dietary protein and natural organic acids such as citric acid improve zinc bioavailability, meaning they increase the percentage of zinc your body absorbs and puts to use. Because the body has no specialized storage system for zinc, maintaining a steady dietary intake helps support healthy zinc levels in the eyes and throughout the rest of the body.

- **Your body carefully regulates zinc every day without you noticing it** – The review explained that only a tiny fraction of your body's approximately 2 to 3 grams of total zinc circulates in the bloodstream at any given time. Most of it is stored inside muscles, bones and other tissues where it performs essential biological functions. Specialized transport proteins in the intestines increase zinc absorption when dietary intake falls, conserving tissue zinc.

When intake is high, however, the review reports that the body compensates mainly by increasing how much zinc it excretes in stool rather than by absorbing less. So, a large supplemental dose is not automatically self-limiting.

At the same time, the digestive tract, liver and other organs continually recycle and redistribute zinc to maintain stable levels. This tightly regulated system helps ensure that delicate retinal cells receive the zinc they need while protecting them from the harmful effects of excess exposure.

- **Researchers discovered that excess zinc released after nerve injury speeds optic nerve cell death** – Research from a completely different corner of eye science reinforces why zinc balance matters so much. Scientists at Boston Children's Hospital and Harvard Medical School were studying retinal nerve damage in [glaucoma](#) when they made an unexpected discovery about zinc itself.^{8,9}

They found that after an acute optic nerve injury in mice, zinc flooded out of neighboring amacrine cells – specialized retinal nerve cells that communicate with retinal ganglion cells – within about one hour of the injury. Surprisingly, the zinc didn't come from the damaged ganglion cells themselves.

Instead, excess zinc spread from surrounding cells and contributed to the death of retinal ganglion cells, the neurons that form the optic nerve and carry visual information from your eye to your brain. In laboratory mice, researchers removed excess zinc using a process called chelation, which binds and removes specific metals.

Ganglion cells survived longer and even regenerated after treatment, and the protective effect remained even when treatment was delayed for several days because zinc toxicity developed gradually instead of immediately.

These findings suggest that maintaining normal zinc balance is just as important as preventing zinc deficiency and raise the question of whether limiting excess zinc could help protect the optic nerve in diseases such as glaucoma. Note that this work has only been demonstrated in laboratory animals so far and has not been tested in people.

Protect Your Eyes by Supporting Them Every Day

Your eyesight depends on much more than a single supplement. Nutrition is one part of a long-term strategy to help your retina stay healthy for as many years as possible.

The research in this article shows that zinc plays an important role in normal retinal function and that a specific combination of nutrients slowed disease progression only in people who already had intermediate AMD or advanced disease in one eye. That makes it even more important to build a strong nutritional foundation before your vision begins to decline.

- 1. Build your meals around foods that naturally provide zinc first** — Your body absorbs zinc from foods such as oysters, grass fed beef, lamb, and other animal foods more efficiently than from many plant sources. If you regularly eat these foods, you give your eyes a steady supply of the mineral they need every day instead of relying

entirely on supplements. It's usually best to meet your nutritional needs through whole foods whenever possible because they provide many nutrients that work together.

Next, understand the participants the studied formula was tested on. The research evaluated one specific combination of antioxidant vitamins and zinc in people who already had a high risk of progressing to advanced AMD. No benefit was observed in people with healthy eyes or early-stage disease.

The formula uses zinc at a dose well above the general upper intake level and was given under clinical supervision, so whether it is appropriate for you is a decision for your eye care provider based on your disease stage — not something to start on your own.

- 2. Avoid thinking that more zinc is always better** — Your retina depends on balanced zinc levels. Too little interferes with normal eye function, while too much also disrupts healthy retinal cells. Stay within the recommended daily intake that was referenced earlier — 11 mg a day for men and 8 mg for women, with an upper limit of 40 mg for adults — unless your eye care provider has specifically recommended the AREDS formula based on your disease stage.

Higher doses aren't automatically better; as the research in this article shows, excess zinc can damage the very cells you're trying to protect. It is also worth knowing that when AREDS2 compared 80 mg of zinc against 25 mg, it found no significant difference in protection.

- 3. Help your body absorb the zinc you eat** — If you eat large amounts of whole grains or legumes, remember that they contain natural compounds called phytates that reduce zinc absorption. Eating zinc-rich animal foods as part of a balanced meal helps improve the amount your body actually uses. Making small adjustments to your daily meals often produces more lasting results than focusing only on pills.

FAQs About Antioxidant Vitamins and Zinc for AMD

Q: Who benefits most from antioxidant vitamins and zinc for AMD?

A: The strongest evidence applies to people who already have intermediate AMD or advanced disease in one eye. A specific combination of antioxidant vitamins and zinc slowed progression to advanced disease and helped preserve vision in this high-risk group. The same benefit wasn't seen in people with no signs of the disease or only early-stage AMD. The formula was studied at doses well above general intake guidance, so whether it fits your situation is a question for your eye care provider.

Q: Why is zinc so important for healthy vision?

A: Zinc supports many of the retina's normal functions, including cell repair, communication between retinal cells and the activity of protective enzymes. Your eyes require a steady supply of zinc to keep these processes working properly, but research also shows that both zinc deficiency and excessive zinc disrupt normal retinal function.

Q: Is taking more zinc better for protecting my eyesight?

A: No. The research consistently shows that balance is more important than taking the highest possible dose. Too little zinc interferes with healthy retinal function, while excessive amounts increase oxidative damage and weaken the eye's natural defense systems. The goal is to maintain healthy zinc levels, not simply consume more.

Q: Can I get enough zinc from food, or do I need a supplement?

A: Many people obtain zinc through foods such as oysters, grass fed beef, lamb and other animal foods, which provide zinc in a form your body absorbs efficiently. Supplements aren't a replacement for a nutrient-rich diet. However, people with intermediate or advanced AMD who fit the criteria studied in the AREDS trial received the greatest benefit from the specific nutrient formula tested in that research.

This article is for informational purposes only and does not constitute medical advice. Consult a qualified health care provider before making changes to your health regimen.

Sources and References

- ^{1, 2} [National Eye Institute October 12, 2001](#)
- ³ [Arch Ophthalmol. 2001 Oct;119\(10\):1417-36](#)
- ⁴ [National Institutes of Health, Zinc](#)
- ⁵ [JAMA 2013 May 15;309\(19\):2005-15](#)
- ^{6, 7} [Progress in Retinal and Eye Research 2025 Sep 19;109:101404](#)
- ⁸ [PNAS. 2017;114\(2\):E209–E218. PMID 28049831](#)
- ⁹ [Glaucoma Research Foundation, May 9, 2022](#)