

Could Your Bedtime Be Ruining Your Morning Poop?

Analysis by [Dr. Joseph Mercola](#)

September 27, 2026

STORY AT-A-GLANCE

- › Your digestive system follows a built-in daily rhythm, so when you go to bed and wake up at consistent times, your body is better prepared for a regular morning bowel movement
- › Poor sleep does more than leave you tired — it disrupts gut bacteria, stress hormones and the communication between your brain and digestive tract, making constipation and other digestive problems more likely
- › Staying up late, working overnight shifts and rushing through your morning all interfere with your body's natural urge to empty your bowels, even if you eat a healthy diet
- › Simple habits such as finishing dinner at least three hours before bedtime, taking a short walk after eating, staying hydrated during the day and using a footstool while on the toilet make bowel movements easier and more comfortable
- › Instead of relying on coffee or laxatives, treating sleep as part of your digestive health gives you another effective way to build more predictable, regular bowel movements over the long term

When a morning bowel movement doesn't happen on schedule, the usual suspect is yesterday's dinner. More often, the real culprit is what happened after dinner — specifically, how you slept and when. Your digestive tract doesn't simply respond to what you eat. It also responds to how well you sleep and, just as importantly, when you sleep.

That connection gets overlooked because conventional advice almost always focuses on fiber, water and exercise. Those recommendations matter, but they leave out a factor that influences your digestion from the moment you close your eyes until the moment your feet hit the floor. Your evening routine, your bedtime consistency and even how much time you allow yourself after waking up all play a role in whether your body cooperates the next morning.

Rather than relying on coffee or hoping for better luck tomorrow, you have far more control over your morning than you might realize. The research below explains exactly how your sleep schedule shapes your digestive rhythm and what you can do each evening to support a better morning.

Your Digestive System Follows a Built-In Daily Schedule

A review published in *Current Treatment Options in Gastroenterology* examined how sleep affects the digestive system in both healthy people and those with disorders of gut-brain interaction.¹ The researchers evaluated evidence covering normal digestive function, sleep disorders and conditions such as [irritable bowel syndrome](#) (IBS), while also reviewing available treatments that target sleep.

Their goal was to determine whether [poor sleep](#) makes digestive symptoms worse, whether digestive problems disrupt sleep or whether both influence each other. The researchers concluded that sleep deserves much more attention in digestive care, noting, "Sleep, a cornerstone of lifestyle management, appears to be the forgotten factor."

- **Your digestive tract intentionally slows down overnight before becoming active again —** Every part of your digestive system follows a predictable daily rhythm. According to the review, saliva production falls by about 50% overnight, stomach emptying slows during the evening, and the waves of muscle contractions that move food through your intestines become much less active while you're asleep.

Even the muscles of your colon reduce their normal activity during the night. These changes don't mean digestion has stopped. Your digestive system is following a programmed rest period, slowing down overnight so it can ramp back up when you wake.

- **Your gut and brain communicate constantly through hormones, nerves and gut bacteria —** The review explained that your digestive tract actually contains up to 400 times more [melatonin](#) than your pineal gland in the brain. Inside your intestines, melatonin helps regulate gut movement and supports healthy gut bacteria.

At the same time, sleep deprivation activates your body's main stress response system, called the hypothalamic-pituitary-adrenal (HPA) axis, increasing stress hormones that weaken the intestinal barrier and interfere with normal gut-brain communication. The

researchers also found that your gut bacteria follow their own daily rhythm, so irregular sleep disrupts the balance of microbes that help digestion stay on track.

- **Poor sleep predicted worse digestive symptoms better than digestive symptoms predicted poor sleep** — People with IBS experienced sleep disturbances far more often than healthy adults, with a pooled prevalence of 37.6%. Many participants actually logged more hours of sleep than healthy individuals but still woke up feeling unrested; their sleep was fragmented by more frequent awakenings and spent disproportionately in lighter stages.

Those repeated awakenings were associated with greater abdominal pain, more digestive distress and lower quality of life. Researchers also found that insomnia predicted worse abdominal pain, anxiety and fatigue the following day, while digestive symptoms didn't consistently predict the following night's sleep quality.

- **Disrupted body clocks increased digestive problems in everyday life** — The review highlighted several real-world examples showing what happens when your **internal clock falls out of sync**. Waking up naturally increases colon contractions that move stool toward the rectum, preparing your body for a morning bowel movement. When sleep schedules constantly change, that timing becomes less predictable.

Researchers also reported that IBS occurred 81% more often among shift workers than among people with regular daytime schedules. Shift work often brings irregular eating habits and greater intake of processed foods, creating multiple pressures on normal digestive function at the same time.

- **Sleep-focused treatments improved digestive health alongside better sleep** — Because sleep influences so many parts of digestion, the researchers also reviewed therapies that target sleep rather than bowel symptoms alone. They found sufficient evidence that melatonin reduced overall IBS symptom severity while improving quality of life.

Cognitive behavioral therapy for insomnia also improved insomnia severity, sleep efficiency and overall sleep quality, with one study showing improvements in both sleep and digestive symptoms among college students with IBS and insomnia.

Based on the evidence reviewed, the researchers concluded that improving sleep deserves a much larger role in first-line digestive care alongside diet, exercise and stress management because healthy digestion depends on healthy sleep from the very beginning.

The research makes a convincing case that sleep shapes digestion at every level — hormones, gut bacteria, colon contractions, even the strength of your intestinal lining. The practical question is what that means for your evening routine and your morning.

Your Evening Routine Determines How Easily You Poop the Next Morning

As noted in a news feature published by EatingWell, a consistent bedtime teaches your digestive system when it's time to go.² Gastroenterologist Dr. Catherine Ngo explains that "consistent, high-quality sleep helps the digestive system maintain predictable patterns of motility, which supports regular bowel habits." Instead of wondering whether your body will cooperate each morning, keeping a regular bedtime teaches your digestive tract when to rest and when to prepare for elimination.

Ngo also explains that going to bed and waking up at roughly the same time each day creates predictability and safety for your nervous system. Because your digestive tract responds to signals from both your brain and your gut, that steady schedule helps your colon anticipate periods of activity and rest, making it easier to move stool forward before you even head to the bathroom.

- **Late nights disrupt the hormones that help keep digestion on schedule** — Gastroenterologist Dr. Rucha Shah explains that staying awake far later than normal throws off the timing of [cortisol](#), your body's primary stress hormone.³ As she puts it, "When people are staying up really late, those cortisol levels are spiking at times that they're not normally used to ... and so then it creates this pattern where digestion gets thrown off."

Shah adds that this problem often appears in people who work overnight shifts because their internal body clock rarely follows a regular daily rhythm. When your sleep schedule constantly changes, your digestive system loses the consistency it depends on.

- **Human research linked poor sleep to measurable changes in the gut microbiome —** A 2026 systematic review analyzed 41 human studies published between 2016 and 2025 that examined how sleep duration, sleep quality, insomnia and disrupted body clocks affect the gut microbiome — the community of bacteria and other microbes living in your digestive tract.⁴

Researchers found that poor sleep and circadian disruption were associated with changes in both the types of gut bacteria present and the beneficial substances they produce, including **short-chain fatty acids**, vitamins, hormones, neurotransmitters and bile acids that help regulate digestion, metabolism and immune function.

While the findings varied between studies because of differences in diet, age, health and research methods, the review concluded that improving sleep and circadian rhythms supports a healthier gut microbiome, giving you another reason to treat a consistent bedtime as part of your digestive health routine instead of simply a way to feel more rested.

- **Rushing through your morning trains your body to ignore its own signals —** The experts point out that many people accidentally work against their body's natural urge to have a bowel movement simply because they're trying to get out the door quickly. Ngo explains that "if rushed in the morning, you don't have the time to relax and let the bowels evacuate."⁵

Shah adds that successful bowel movements require the muscles around the rectum to relax. If you repeatedly suppress the urge because you're short on time, your body gradually becomes less responsive to those natural signals, making regular bowel movements more difficult over time.

- **A simple change in body position helps stool pass more easily —** One practical recommendation involves placing a small stool beneath your feet while sitting on the toilet. According to Shah, raising your knees above your hips helps open the muscles involved in **bowel movements**, making it "exponentially easier to poop."

It's a straightforward mechanical adjustment that changes the angle of your lower bowel, helping stool move through more comfortably without medication, supplements or expensive equipment.

Make Your Evening Routine Work for Your Gut

If your mornings rarely go as planned, don't focus only on breakfast or coffee. Start the night before. Your digestive system follows a daily rhythm, and the habits you repeat every evening shape what happens the next morning. Try making a few simple changes and sticking with them long enough for your body to recognize the pattern. Consistency matters far more than perfection.

- 1. Keep the same bedtime every day** — Your digestive tract performs best when your sleep schedule stays predictable. Go to bed and wake up at roughly the same time every day, including weekends. That steady routine helps your body recognize when it's time to rest and when it's time to prepare for a morning bowel movement. If you stay up late several nights in a row, don't expect your digestive system to stay perfectly regular. Your body responds best to repetition.
- 2. Give yourself enough time after you wake up** — If you usually race through your morning, build in an extra 15 to 30 minutes. Your digestive system works better when you aren't under pressure. Don't ignore the urge to have a bowel movement because you're running late. Repeatedly holding it trains your body to become less responsive over time, making [constipation](#) more likely.

If you often feel rushed, shifting your bedtime earlier is one of the simplest ways to create more time the next morning.

- 3. Support digestion after dinner instead of collapsing on the couch** — Finish eating at least three hours before bedtime whenever possible so your digestive system has time to process your meal before you fall asleep. A light walk after your evening meal helps keep food moving through your digestive tract.

Even 10 to 20 minutes is enough to encourage normal stomach emptying and healthy intestinal movement. Pair that habit with steady hydration throughout the day instead of drinking large amounts of water right before bed, which interrupts sleep with nighttime bathroom trips.

- 4. Use better bathroom mechanics** — Your body was designed to empty your bowels in more of a squat than a seated position. Place a small footstool under your feet so your knees sit higher than your hips while you're on the toilet. That simple adjustment straightens the lower

part of your bowel and allows the muscles involved in a bowel movement to relax more completely. If you haven't tried it, you'll often notice the difference immediately.

- 5. Build regularity through daily habits instead of quick fixes** — Your digestive system responds to patterns, not shortcuts. Protect your sleep as carefully as you protect your diet. If your gut tolerates it well, gradually increase **fiber-rich whole foods** and stay physically active every day, but recognize that without consistent sleep, those habits lose some of their power.

I also encourage you to keep a simple journal for two weeks that tracks your bedtime, wake time and bowel habits. You'll often spot patterns that weren't obvious before, making it much easier to identify the routines that keep your digestive system working on schedule.

FAQs About Sleep and Bowel Movements

Q: Does poor sleep really affect my bowel movements?

A: Yes. Your digestive system follows a daily rhythm that depends on consistent sleep. Irregular bedtimes, poor-quality sleep and disrupted body clocks interfere with the normal patterns that move stool through your colon, making it harder for your body to stay regular.

Q: Why are morning bowel movements more common than bowel movements later in the day?

A: Your colon naturally becomes more active after you wake up, helping move stool toward the rectum. A regular sleep schedule reinforces this built-in timing, while inconsistent bedtimes make the process less predictable and increase the likelihood of constipation or delayed bowel movements.

Q: What nighttime habits help promote a morning bowel movement?

A: Going to bed at the same time each night, finishing your last meal at least three hours before bedtime, taking a 10- to 20-minute walk after dinner and staying well hydrated throughout the day all help support your digestive system's natural rhythm. These simple

habits work together to encourage more regular bowel movements.

Q: Why does rushing in the morning make it harder to poop?

A: Your body needs time to respond to its natural urge to have a bowel movement. If you repeatedly ignore that urge because you're in a hurry, your body gradually becomes less responsive to those signals. Giving yourself an extra 15 to 30 minutes each morning makes it easier for your bowels to empty completely.

Q: Is there a simple way to make bowel movements easier without medication?

A: Yes. Placing a small footstool under your feet while sitting on the toilet raises your knees above your hips, creating a more natural squatting position. This helps relax the muscles involved in bowel movements and allows stool to pass more comfortably, especially when combined with healthy sleep habits and a consistent daily routine.

This article is for informational purposes only and does not constitute medical advice. Consult a qualified healthcare provider before making changes to your health regimen.

Test Your Knowledge with Today's Quiz!

Take today's quiz to see how much you've learned from [yesterday's Mercola.com article](#).

What can cause tonsil stones to keep coming back?

[Bacteria and debris collecting in tonsil folds](#)

[Drinking too much water, causing debris to become stuck](#)

[Eating crunchy fruits and vegetables](#)

[Breathing only through the nose](#)

Sources and References

- ¹ Current Treatment Options in Gastroenterology October 10, 2023, Volume 21, Pages 435-446
- ^{2, 3, 5} EatingWell July 7, 2026
- ⁴ Sleep Medicine Reviews April 2026, Volume 86, 102256