Referencias:

[1] Journal of Alzheimer's Disease March 9, 2012

[2] Nat Med. 2012 Jan 29;18(2):291-5.

[3] Neuroimage. 2011 Mar 1;55(1):32-8. Epub 2010 Nov 30.

[4] WSJ.com January 31, 2012

[5] Green tea or rosemary extract added to foods reduces nonheme-iron absorption. Am J Clin Nutr. 2001 Mar;73(3):607-12. PMID: 11237939

[6] Iron chelation in the biological activity of curcumin. Free Radic Biol Med. 2006 Apr 1;40(7):1152-60. PMID: 16545682

[7] GreenMedInfo.com, Astaxanthin Research Page

[8] Astaxanthin and peridinin inhibit oxidative damage in Fe(2+)-loaded liposomes: scavenging oxyradicals or changing membrane permeability? Biochem Biophys Res Commun. 2001 Oct 19;288(1):225-32. PMID: 11594777

[9] 2011 Alzheimer's Diseases Facts and Figures (PDF)

[10] Is Vitamin D Important for Preserving Cognition? A Positive Correlation of Serum 25-Hydroxyvitamin D Concentration with Cognitive Function, Archives of Biochemistry and Biophysics , April 15, 2007: 460(2); 202-205, Robert J. Przybelski and Neil C. Binkley.

[11] Beauty and the Beast: B12, Homocysteine, and the Brain: A Bemusing Saga!, Neurology , October 19, 2010: 75(16); 1402-3, S. Seshadri.

[12] Vitamin B12 and Folate in Relation to the Development of Alzheimer's Disease, Neurology , May 8, 2001: 56(9); 1188-1194, H-X. Wang, BA, et al.

[13] A Diet Enriched with the Omega-3 Fatty Acid Docosahexaenoic Acid Reduces Amyloid Burden in an Aged Alzheimer Mouse Model, Journal of Neuroscience , March 23, 2005: 25(12); 3032-40, G. P. Lim, et al.

[14] Voluntary Exercise Decreases Amyloid Load in a Transgenic Model of Alzheimer's Disease, Journal of Neuroscience , April 27, 2005: 25(17); 4217-4221, Paul A. Adlard, et al.

[15] PPARγ Co-activator-1α (PGC-1α) Reduces Amyloid-β Generation through a PPARγ-Dependent Mechanism, Journal of Alzheimer's Disease , 2011: 25(1); 151-62, L. Katsouri, et al.

[16] Use of Anticholinergics and the Risk of Cognitive Impairment in an African American Population, Neurology , July 13, 2010: 75(2); 152-9, N. L. Campbell, et al.